

PATIENT INFORMATION				
Patient Name:		Caregiver (if applicable):		
Address:		City:	State:	Zip:
Main Phone:	Alternate Phone:		Email:	Last 4 of SSN (required):
Date of Birth:	Male or Female	Height (required): _____ cm or inches	Weight (required): _____ lb or kg	
CLINICAL INFORMATION				
Primary ICD-10 Code:	Is patient currently on therapy? Yes No		Previous tried/failed therapies:	Allergies:
REQUIRED (ALL): Has the patient tested NEGATIVE for TB? Yes No Pending Result:_____ Test Date: _____				
REQUIRED (Infliximab, Simponi Aria, Tremfya): Has the patient tested NEGATIVE for Hep B infection? Yes No Pending Result:_____ Test Date: _____				
Please include face sheet, copy of insurance cards (front and back), H&P, clinicals and any available lab results.				
PRESCRIPTION AND ORDERS				
Medication	Dosing		Quantity	Refills
Infliximab (Remicade, Inflectra, Renflexis, Avsola*)	Induction: 5 mg/kg IV at 0, 2 & 6 weeks Other: _____	Maintenance: ____mg/kg IV every 8 weeks after induction Other: _____	3 doses for induction 8-week supply for maintenance Other: _____	x 1 year unless not ed otherwise: ____refills
	*Pharmacy will select infliximab product unless preferred brand is indicated. Preferred brand: _____			
Simponi Aria®	Induction: Adults: 2 mg/kg IV at 0 & 4 weeks Other: _____	Maintenance: Adults: 2 mg/kg IV every 8 weeks after induction Other: _____	2 doses for induction 8-week supply for maintenance Other: _____	x 1 year unless not ed otherwise: ____refills
Stelara® (Crohn's disease/ Ulcerative Colitis)	Induction: Adults: ≤55 kg: 260 mg IV at week 0 Adults: >55 kg - 85 kg: 390 mg IV at week 0 Adults: >85 kg: 520 mg IV at week 0	Maintenance: Adults: 90 mg SUBQ every 8 weeks after induction Other: _____	one dose for induction 8-week supply for maintenance	x 1 year unless not ed otherwise: ____refills
Stelara® (Plaque Psoriasis and Psoriatic Arthritis)	Induction: Adults: ≤100 kg: 45 mg SUBQ at 0 & 4 weeks Adults: >100 kg: 90 mg SUBQ at 0 & 4 weeks Other: _____	Maintenance: Adults: ≤100 kg: 45 mg SUBQ every 12 weeks Adults: >100 kg: 90 mg SUBQ every 12 weeks Other: _____	one dose for induction 12-week supply for maintenance	x 1 year unless not ed otherwise: ____refills
Tremfya® (Plaque Psoriasis and Psoriatic Arthritis)	Induction: Adults: 100 mg SUBQ at 0 & 4 weeks	Maintenance: Adults: 100 mg SUBQ every 8 weeks Other: _____	2 doses for induction, no refills 8-week supply for maintenance Other: _____	x 1 year unless not ed otherwise: ____refills
Tremfya® (Ulcerative Colitis)	Induction: Adults: 200 mg IV at 0, 4 & 8 weeks Other: _____	Maintenance: Adults: 100 mg SUBQ at week 16, then every 8 weeks OR Adults: 200 mg SUBQ at week 12, then every 4 weeks	3 doses for induction, no refills 8-week supply for maintenance 4-week supply for maintenance	x 1 year unless not ed otherwise: ____refills
Tremfya® (Crohn's Disease)	Induction: Adults: 200 mg IV at 0, 4 & 8 weeks Adults: 400 mg SUBQ at 0, 4 & 8 weeks	Maintenance: Adults: 100 mg SUBQ at week 16, then every 8 weeks OR Adults: 200 mg SUBQ at week 12, then every 4 weeks	3 doses for induction, no refills 8-week supply for maintenance 4-week supply for maintenance	x 1 year unless not ed otherwise: ____refills
Medication Specific Infusion and Administration: Infusion rate titrated based on the individual product and patient tolerability. Clinical Pharmacist to adjust rates as clinically indicated. Nursing refer to prescription label for patient specific infusion rates.				
Labs:				
Flush Orders (IV infusions ONLY): Flush with 0.9% sodium chloride 5-10 mL pre and post medication and as needed, max per dispense x 60 syringes (refill x 1 year). For CVADs ONLY, may add Heparin 3-5 mL as final flush to maintain patency as needed; PORT 100 units/mL; PICC 10 units/mL, max per dispense x 30 syringes (refill x 1 year). Supplies: Dispense ancillary supplies, including ambulatory infusion pump (IV infusions ONLY), as needed to provide home infusion therapy.				
Premedication 30 minutes prior to infusion: Acetaminophen 500-1000 mg PO, dispense x 1 bottle (refill x 1 year) Diphenhydramine 25-50 mg PO, dispense x 1 bottle (refill x 1 year)		Adverse/Anaphylactic Reactions, per protocol (Subcutaneous): • Epinephrine autoinjector 0.3 mg IM as directed PRN for anaphylaxis, dispense 2pk x 1 (refill x 1 year)		Adverse/Anaphylactic Reactions, per protocol (Intravenous): dispense ana-kit x 1 (refill x 1 year) • Epinephrine 1:1000, 1 mL vial x 1 • Diphenhydramine 50 mg/1 mL vial x 1 • Diphenhydramine 50 mg tab/cap x 1 • 0.9% Sodium Chloride 500 mL bag x 1
Nursing Orders for Home Infusion: Provide skilled nursing for home medication administration as prescribed and per CSI nursing policies. Insert, access, and manage vascular access devices. Deliver nursing care for therapy duration and educate patient on medication, disease state, adverse reactions, and administration. Monitor patient response and report adverse events.				
PRESCRIBER INFORMATION				
Prescriber Name:		Practice site name:		
Address:		City:	State:	Zip:
NPI:	Phone:	Fax:		
Office contact and special instructions:				
Prescriber Authorization: I authorize the pharmacy and its representatives to act as my authorized agent to secure coverage and initiate the insurance prior authorization process for my patient(s), and to sign any necessary forms on my behalf as my authorized agent, including the receipt of any required prior authorization forms and the receipt and submission of patient lab values and other patient data. In the event that this pharmacy determines that it is unable to fulfill this prescription, I further authorize this pharmacy to forward this information and any related materials related to coverage of the product to another pharmacy of the patient's choice or in the patient's insurer's provider network.				
Prescriber's Signature:			Date:	